
The Higgs sector in the complex MSSM: QCD corrections

Heidi Rzehak

Paul-Scherrer-Institut

in coll. with S. Heinemeyer, W. Hollik and G. Weiglein

Outline

- ▶ Higgs bosons in the complex MSSM
- ▶ Mass of the lightest Higgs boson
- ▶ Higher order contributions to this mass

Higgs bosons

At Born level: no CP-violation:

- ▶ one phase in the Higgs potential: $V_{\text{Higgs}} = \dots + \epsilon_{ij} |m_3^2| e^{i\varphi} m_3^2 H_1^i H_2^j + \dots$
elimination via Peccei-Quinn transformation
- ▶ phase difference of Higgs doublets:
vanishes because of minimum condition

Physical mass eigenstates (at Born level):

- ▶ 5 Higgs bosons: 3 neutral H^0, h^0, A^0 ; 2 charged H^\pm

Masses of the Higgs bosons:

- ▶ not all independent: here: H^\pm -mass M_{H^\pm} (and $\tan \beta$) as free parameter
 $\tan \beta = \frac{v_2}{v_1}$: ratio of the Higgs vac. expect. values
- ▶ lightest Higgs boson: h^0


Mass of the lightest Higgs boson

Upper theoretical Born mass bound: $M_{h^0} \leq M_Z = 91 \text{ GeV}$

with quantum corrections of higher orders: $M_{h^0} \lesssim 135 \text{ GeV}$

dependent on the MSSM parameters:
particularly on parameter phases

- Discovery of the Higgs boson:




- Before the discovery: Exclusion of parts of the parameter space

Determination of the Higgs masses

Two-point-function:

$$\Gamma(k^2) = k^2 - M_{\text{Born}}^2 + \begin{pmatrix} \hat{\Sigma}_{H^0 H^0}(k^2) & \hat{\Sigma}_{H^0 h^0}(k^2) & \hat{\Sigma}_{H^0 A^0}(k^2) \\ \hat{\Sigma}_{H^0 h^0}(k^2) & \hat{\Sigma}_{h^0 h^0}(k^2) & \hat{\Sigma}_{h^0 A^0}(k^2) \\ \hat{\Sigma}_{H^0 A^0}(k^2) & \hat{\Sigma}_{h^0 A^0}(k^2) & \hat{\Sigma}_{A^0 A^0}(k^2) \end{pmatrix}$$

\uparrow
 diagonal matrix with squared Born masses
 $\text{diag}(M_{H_{\text{Born}}^0}^2, M_{h_{\text{Born}}^0}^2, M_{A_{\text{Born}}^0}^2)$



determining the zero of $\det(\Gamma(k^2)) \Rightarrow M_{h_1}, M_{h_2}, M_{h_3}$

Real parameters:


$$\hat{\Sigma}_{H^0 A^0}(k^2) = \hat{\Sigma}_{h^0 A^0}(k^2) = 0 \Rightarrow M_{h^0} = M_{h_1}$$

no mixing between CP-even and CP-odd states

Renormalized two-loop self energies

Calculation of the dominant two-loop contributions ($\alpha_t = \lambda_t^2/(4\pi)$):

↑
Yukawa coupling



- ▶ Terms of order $\mathcal{O}(\alpha_t \alpha_s)$ for complex parameters
- ↖ new

known: two-loop leading-log contributions [Pilaftsis, Wagner]

[Carena, Ellis, Pilaftsis, Wagner]


- ▶ Extraction of the relevant terms (equiv. to eff. potential approach):
 - use vanishing external momenta $\hat{\Sigma}^{(2)}(0)$
 - use vanishing electroweak gauge couplings g, g'

Renormalized two-loop self energies

Calculation of the dominant two-loop contributions :

▶ within an **on-shell** scheme in the Higgs sector:

- no shift of the minimum of the Higgs potential
- define the H^\pm -mass M_{H^\pm} as the **pole mass**
 \Rightarrow directly related to a **physical observable**



▶ with parameters of the top (bottom) sector defined at one-loop:

- top quark mass and top squark masses on-shell
- generalization of the mixing angle condition:

$$\widetilde{\text{Re}}\hat{\Sigma}_{\tilde{t}_{12}}(m_{\tilde{t}_1}^2) + \widetilde{\text{Re}}\hat{\Sigma}_{\tilde{t}_{12}}(m_{\tilde{t}_2}^2) = 0$$

Phases in couplings

Phases relevant at two-loop level:

► squark sector:

- phase φ_{A_t} of the trilinear coupling A_t
- phase of μ (small), μ : Higgsino mass parameter


constraints from
measurements of
electr. dipole moments

soft breaking
parameter

► gluino sector:



- phase $\varphi_{\tilde{g}}$ of the gluino mass parameter
in the gluino-squark-quark-vertex

Results: φ_{A_t} -dependence (small M_{H^\pm})




- Quantum corrections of $\mathcal{O}(\alpha_t \alpha_s)$ change the qualitative behaviour of M_{h_1} .

Results: φ_{A_t} -dependence (large M_{H^\pm})



- Quantum corrections of $\mathcal{O}(\alpha_t \alpha_s)$ change the qualitative behaviour of M_{h_1} .
- Qualitative behaviour of M_{h_1} depends strongly on $|A_t|$.



Results: $|A_t|$ -dependence (large M_{H^\pm})



- One-loop: $\varphi_{A_t} = \pi$ “shifts” towards lower values of $|A_t|$ with respect to $\varphi_{A_t} = 0$.
- Two-loop: also the value of the maximum of M_{h_1} depends on the phases.


Results: φ_{A_t} - versus φ_{X_t} -dependence (large M_{H^\pm})

size of the squark mixing: $X_t := A_t - \mu^* \cot \beta$



- Quantum corrections are smaller for constant absolute value of the squark mixing, $|X_t| = \text{const.}$


Results: $\varphi_{\tilde{g}}$ -dependence



$\tan\beta = 10, M_{H^\pm} = 500$ GeV ——— $\tan\beta = 10, M_{H^\pm} = 150$ GeV ●
 $\tan\beta = 5, M_{H^\pm} = 500$ GeV - - - x - - - $\tan\beta = 5, M_{H^\pm} = 150$ GeV ▲
 $\tan\beta = 3, M_{H^\pm} = 500$ GeV ····· $\tan\beta = 3, M_{H^\pm} = 150$ GeV ○

- M_{h_1} depends rather weakly on the phase $\varphi_{\tilde{g}}$.

Results: dependence on the gluino mass $m_{\tilde{g}}$



- Large effects in the threshold region: $m_{\tilde{t}_2} = m_{\tilde{g}} + m_t$

Conclusions

- ▶ Quantum corrections are important for a precise prediction of the mass of the lightest Higgs boson M_{h_1} :
 - They can induce CP-violation.
 - Dominant corrections: from the top sector
- ▶ Contributions of $\mathcal{O}(\alpha_t \alpha_s)$ for complex parameters:
 - Phases are relevant at the two-loop level.
 - Two-loop contributions can change qualitative behaviour of M_{h_1} .
 - are currently included into FeynHiggs.